

Quasi-local formulation of the mirror TBA

János Balog and Árpád Hegedűs

*Research Institute for Particle and Nuclear Physics, Hungarian Academy of Sciences,
H-1525 Budapest 114, P.O.B. 49, Hungary*

ABSTRACT: We present a method of removing all infinite sums from the various forms of the mirror TBA equations and the energy formula of the AdS/CFT spectral problem. This new formulation of the TBA system is quasi-local because Y -functions that are connected by the TBA equations are at most next to nearest neighbors with respect to the Y -system diagram of AdS/CFT.

Contents

1. Introduction	1
2. TBA sums	3
3. Energy formula and asymptotic form of the identity	6
4. The quasi-local TBA equations	8
5. Conclusion	10

1. Introduction

One of the most important problems in testing the AdS/CFT correspondence [1] in the planar limit is to determine the finite size spectrum of the $AdS_5 \times S^5$ superstring sigma model [2]. After integrability was discovered in the string worldsheet theory, the mirror Thermodynamic Bethe Ansatz (TBA) technique was proposed [3, 4] to determine nonperturbatively the spectrum of the string theory. The TBA equations for AdS/CFT were first derived for the ground state [5, 6, 8, 7, 9] and then using an analytic continuation trick [10] TBA equations were conjectured for certain classes of states in the $\mathfrak{sl}(2)$ sector of the theory [11, 12, 13].

An important property of the TBA equations is that the unknown functions (Y -functions) satisfy the so called Y -system functional relations [14]. Recently an alternative derivation of the TBA equations appeared in [15] where it has been shown that the complicated ground state TBA equations can be derived from the Y -system if it is supplemented by discontinuity functional equations relating the square-root discontinuities of the various Y -functions. Later the derivation has been extended to excited states [16], too.

The main advantage of the Y -system based approach to the TBA problem is that it gives a deeper insight into the analytic properties of the Y -functions and opens the way to understand the analytic properties of the T and Q functions, which are the elementary building blocks of the underlying Y -system. This knowledge is indispensable if we want to construct an NLIE system formulated in terms of finitely many unknown functions for the AdS/CFT spectral problem.

So far the proposed TBA equations satisfied all tests in both the weak and strong coupling limits. Their correctness has been nicely demonstrated by the convincing agreement with gauge theory results [17, 18, 19, 20] in the weak coupling limit [21, 22, 13] through the generalized Lüscher approach [23, 24, 25, 26, 27], and with string theory results in the strong coupling limit [11, 28, 29, 30, 31].

In spite of the success of TBA technique in AdS/CFT, it has some obvious disadvantages as well. First of all, like all the TBA equations of known sigma-models it contains infinitely many unknown functions, which makes the study of their properties, both analytically and numerically, difficult. Secondly, contrary to the case of relativistic models, these TBA equations cannot be formulated in a local form, namely a Y -function associated to a node of the Y -diagram of the model cannot be expressed, in general, in terms of the Y -functions belonging to the neighboring nodes only.

This non-locality is related to the discontinuity structure of the Y -functions and the discontinuity relations [15]. As a consequence of non-locality all formulations¹ of the mirror TBA equations presented so far contains infinite sums of certain Y combinations convoluted with appropriate kernels on the right hand side of the equations. If expressed in terms of Y -functions, the TBA energy formula has the same property.

The purpose of this paper is to work out a method that enables us to replace all the infinite sums appearing in the TBA approach by expressions containing only a few terms and depending on a few Y -functions only.

Here we focus our attention to the hybrid formulation of the equations [12]. In this case all the infinite sums which appear in the TBA description (including the energy formula) are of the form:

$$\sum_{Q=1}^{\infty} L_Q \star \mathcal{K}_Q, \quad (1.1)$$

where $L_Q = \ln(1 + Y_Q)$ and $\mathcal{K}_Q \equiv \mathcal{K}_Q(u, v)$ is a kernel satisfying the relation:

$$\mathcal{K}_Q - s \star (\mathcal{K}_{Q-1} + \mathcal{K}_{Q+1}) = \delta \mathcal{K}_Q, \quad \mathcal{K}_0 \equiv 0, \quad Q = 1, 2, \dots \quad (1.2)$$

The fact that enables us to transform (1.1) into a finite sum is that for the $\mathcal{K}_Q$ kernels appearing in the TBA equations $\delta \mathcal{K}_Q = 0$, with the exception of a few values of the index Q only. See formulae (4.18)-(4.19).

In this paper we will derive finite expressions for the infinite sums

$$\Omega = \sum_{Q=2}^{\infty} L_Q \star \mathcal{K}_Q. \quad (1.3)$$

The simplification of Ω consists of two main steps. First we extend the definition of the convolution for functions of two variables in a natural way. Next with the help of the simplified TBA equations we can derive a linear integral equation for Ω , which can be solved in Fourier space giving our final formula (2.33).

If we replace the infinite sums of type (1.3) with our new finite formulas (2.33) in the TBA equations and in the energy formula, the resulting formulation of the TBA equations becomes quasi-local² as well. We think that such a quasi-local formulation of the TBA equations not only makes it possible to perform more accurate numerical simulations but

¹Using the terminology of [12] canonical, simplified and hybrid versions of the TBA equations.

²Here quasi-locality means that at most next to nearest neighbor Y -functions are connected by the TBA equations.

it is also an important step towards the NLIE formulation of the planar AdS/CFT spectral problem.

The paper is organized as follows: Section 2 contains the derivation of the finite expression for Ω . In section 3 we discuss the energy formula and evaluate the finite formula for Ω when all Y -functions are replaced by their asymptotic counterparts. In section 4 we discuss the kernels appearing the TBA equations and present the complete set of quasi-local mirror TBA equations. The paper is closed by our conclusions.

Throughout the paper we use the notations, conventions and definitions of [12]. In this short paper we cannot reproduce all the formulae giving the definition of the various kernels appearing in the TBA equations and thus unfortunately the paper is not self-contained. The necessary definitions together with the list of the TBA equations in the form we are using them in this paper can also be found in [16].

2. TBA sums

Since the convolution operation between functions plays a central role in our considerations here we recall the relevant definitions. The objects appearing in these formulas are either “functions” (typically the logarithm of some Y -function) or “kernels”. “Functions” depend on one variable, kernels on two. Some kernels only depend on the difference of the two variables. This is the case for the most important kernel $s(u, v)$:

$$s(u, v) = s(u - v), \quad s(u) = \frac{g}{4 \cosh \frac{\pi g u}{2}}. \quad (2.1)$$

For the “function” $\star$ “kernel” and “kernel” $\star$ “function” type convolutions we have

$$(f \star F)(v) = \int_{-\infty}^{\infty} du f(u) F(u, v), \quad (F \star f)(v) = \int_{-\infty}^{\infty} du F(v, u) f(u) \quad (2.2)$$

and for “kernel” $\star$ “kernel” type convolutions

$$(F_1 \star F_2)(u, v) = \int_{-\infty}^{\infty} dw F_1(u, w) F_2(w, v). \quad (2.3)$$

The modified convolutions

$$(f \hat{\star} F)(v) = \int_{-2}^2 du f(u) F(u, v), \quad (f \check{\star} F)(v) = \left(\int_2^{\infty} + \int_{-\infty}^{-2} \right) du f(u) F(u, v) \quad (2.4)$$

(and similarly for $F_1 \hat{\star} F_2$, $F_1 \check{\star} F_2$) are also used in some of the TBA integral equations.

Note that the standard convolution³ definition between two functions of one variable

$$(f \star^{\text{old}} g)(u) = \int_{-\infty}^{\infty} dw f(u - w) g(w) \quad (2.5)$$

is not always the same as our convolution, but

$$f \star s = f \star^{\text{old}} s, \quad (2.6)$$

³As is well known, $f \star^{\text{old}} g = g \star^{\text{old}} f$ and $\star^{\text{old}}$ reduces to ordinary product after Fourier transformation.

because s is even.

In this paper we will use a new convolution definition, which always maps two functions of two variables into a function of two variables. Functions of one variable will be treated as functions of two variables by writing $f(u, v) \rightarrow f(u - v)$:

$$(F_1 \star F_2)(u, v) = \int_{-\infty}^{\infty} dw F_1(u, w) F_2(w, v), \quad (2.7)$$

$$(f \star F)(u, v) = \int_{-\infty}^{\infty} dw f(u - w) F(w, v), \quad (2.8)$$

$$(f \star g)(u, v) = \int_{-\infty}^{\infty} dw f(u - w) g(w - v) = (f \star^{\text{old}} g)(u - v). \quad (2.9)$$

The new convolution $\star$ is an (in general) non-commutative, associative product of functions.

Finally we note that the prototype TBA equation

$$L = \Lambda \star s, \quad (2.10)$$

where L and Λ are “functions”, can be equivalently written

$$\hat{L} = \hat{\Lambda} \star s, \quad (2.11)$$

where

$$\hat{f}(u) = f(-u) \quad (2.12)$$

for any “function”.

The kernel identity (1.2) can be rewritten as

$$\mathcal{K}_Q - s \star (\mathcal{K}_{Q-1} + \mathcal{K}_{Q+1}) = \delta \mathcal{K}_Q, \quad \mathcal{K}_0 \equiv 0, \quad Q = 1, 2, \dots \quad (2.13)$$

and we also introduce the notation $\bar{\mathcal{K}}_m = s \star \mathcal{K}_m$. $\bar{\mathcal{K}}_m$ satisfies an identity similar to (2.13) with $\delta \bar{\mathcal{K}}_m = s \star \delta \mathcal{K}_m$.

In this section we will present the simplification of the TBA sums working, for simplicity, in the $\mathfrak{sl}(2)$ sector of the theory⁴. We will consider the infinite sums

$$\mathcal{A} = \sum_{Q=2}^{\infty} \hat{L}_Q \star \mathcal{K}_Q \quad (2.14)$$

and

$$\mathcal{B} = \sum_{m=1}^{\infty} \hat{\mathcal{L}}_m \star \bar{\mathcal{K}}_{m+1}. \quad (2.15)$$

After having computed $\mathcal{A}$ we can write our final result as $\Omega(v) = \mathcal{A}(0, v)$.

We recall that $L_Q = \ln(1 + Y_Q)$ and (see ref. [16])

$$\mathcal{L}_m = \ln \left(\tau_m \tau_{m+2} \left(1 + \frac{1}{Y_{m|vw}} \right) \right), \quad m = 1, 2, \dots \quad (2.16)$$

⁴Performing the calculation for the general case is straightforward.

further $r_m = \ln(1 + Y_{m|vw})$ $m = 1, 2, \dots$, $r_0 = 0$ and we define analogously

$$\mathcal{R}_Q = \ln \left(\tau_Q^2 \left(1 + \frac{1}{Y_Q} \right) \right), \quad Q = 2, 3, \dots, \quad \mathcal{R}_1 = \ln \left(1 + \frac{1}{Y_1} \right). \quad (2.17)$$

Using the notation introduced above we write the simplified TBA integral equations (4.1) and (4.2) as [16]:

$$\mathcal{L}_m = r_m + L_{m+1} \star s - (r_{m+1} + r_{m-1}) \star s - \mathfrak{f}_2 \delta_{m,1} \quad m = 1, 2, \dots \quad (2.18)$$

and equivalently

$$\hat{\mathcal{L}}_m = \hat{r}_m + \hat{L}_{m+1} \star s - (\hat{r}_{m+1} + \hat{r}_{m-1}) \star s - \hat{\mathfrak{f}}_2 \delta_{m,1} \quad m = 1, 2, \dots \quad (2.19)$$

Here

$$\mathfrak{f}_2 = H \star s, \quad H = \ln \left(\frac{1 - Y_-}{1 - Y_+} \right). \quad (2.20)$$

Similarly from (4.5)

$$\hat{L}_Q = \hat{\mathcal{R}}_Q + 2\hat{\mathcal{L}}_{Q-1} \star s - (\hat{\mathcal{R}}_{Q+1} + \hat{\mathcal{R}}_{Q-1}) \star s - \hat{\mathfrak{f}}_1 \delta_{Q,2} \quad Q = 2, 3, \dots \quad (2.21)$$

where

$$\mathfrak{f}_1 = \ln \tau_1^2 \star s. \quad (2.22)$$

Now we take the convolution of (2.21) with $\mathcal{K}_Q$ and sum over Q from 2 to infinity. After rearranging terms and using the kernel identity (2.13) we get

$$\mathcal{A} = 2\mathcal{B} + \sum_{Q=2}^{\infty} \hat{\mathcal{R}}_Q \star \delta \mathcal{K}_Q + \hat{\mathcal{R}}_2 \star \bar{\mathcal{K}}_1 - \hat{\mathcal{R}}_1 \star \bar{\mathcal{K}}_2 - \hat{\mathfrak{f}}_1 \star \mathcal{K}_2. \quad (2.23)$$

After performing similar transformations we get from (2.19)

$$\mathcal{B} = \sum_{m=1}^{\infty} \hat{r}_m \star \delta \bar{\mathcal{K}}_{m+1} + \hat{r}_1 \star \bar{\bar{\mathcal{K}}}_1 + s \star s \star \mathcal{A} - \hat{\mathfrak{f}}_2 \star \bar{\mathcal{K}}_2. \quad (2.24)$$

Eliminating $\mathcal{B}$ from (2.23) and (2.24) we get a linear integral equation of the form

$$\mathcal{A} - 2s \star s \star \mathcal{A} = \omega \quad (2.25)$$

with

$$\begin{aligned} \omega = & \sum_{Q=2}^{\infty} \hat{\mathcal{R}}_Q \star \delta \mathcal{K}_Q + \hat{\mathcal{R}}_2 \star \bar{\mathcal{K}}_1 - \hat{\mathcal{R}}_1 \star \bar{\mathcal{K}}_2 - \hat{\mathfrak{f}}_1 \star \mathcal{K}_2 \\ & + 2 \sum_{m=1}^{\infty} \hat{r}_m \star \delta \bar{\mathcal{K}}_{m+1} + 2\hat{r}_1 \star \bar{\bar{\mathcal{K}}}_1 - 2\hat{\mathfrak{f}}_2 \star \bar{\mathcal{K}}_2. \end{aligned} \quad (2.26)$$

The linear equation (2.25) can easily be solved since the second argument of the unknown $\mathcal{A}(u, v)$ is a “spectator” only and the essential part of the problem can be formulated in

terms of kernels depending on differences only and it can be algebraically solved in Fourier space. The solution is of the form

$$\mathcal{A} = M \star \omega \quad (2.27)$$

with

$$M = \delta + 2s_{1/2}, \quad s_{1/2}(u) = \frac{1}{2}s\left(\frac{u}{2}\right), \quad (2.28)$$

where δ is the Dirac delta function. This solution further satisfies

$$M \star s = \sigma_{1/2}, \quad \sigma_{1/2} = s_{1/2}^+ + s_{1/2}^-, \quad \sigma_{1/2}(u) = \frac{g}{2\sqrt{2}} \frac{\cosh \frac{\pi g u}{4}}{\cosh \frac{\pi g u}{2}}. \quad (2.29)$$

A further useful identity is

$$M \star s \star s = s_{1/2}. \quad (2.30)$$

Putting everything together, $\mathcal{A}$ can be expressed explicitly:

$$\begin{aligned} \mathcal{A} = & \sum_{Q=2}^{\infty} \hat{\mathcal{R}}_Q \star (\delta \mathcal{K}_Q + 2s_{1/2} \star \delta \mathcal{K}_Q) + \hat{\mathcal{R}}_2 \star \sigma_{1/2} \star \mathcal{K}_1 - \hat{\mathcal{R}}_1 \star \sigma_{1/2} \star \mathcal{K}_2 \\ & - \ln \hat{\tau}_1^2 \star \sigma_{1/2} \star \mathcal{K}_2 + 2 \sum_{m=1}^{\infty} \hat{r}_m \star \sigma_{1/2} \star \delta \mathcal{K}_{m+1} \\ & + 2\hat{r}_1 \star s_{1/2} \star \mathcal{K}_1 - 2\hat{f}_2 \star \sigma_{1/2} \star \mathcal{K}_2. \end{aligned} \quad (2.31)$$

In all applications we need in this paper the kernel functions satisfy

$$\delta \mathcal{K}_Q = 0, \quad Q \geq 3 \quad (2.32)$$

and furthermore we are interested in Ω only and thus the original convolutions can be used. Our final result is

$$\begin{aligned} \Omega = & \mathcal{R}_2 \star (\delta \mathcal{K}_2 + 2s_{1/2} \star \delta \mathcal{K}_2) + \mathcal{R}_2 \star \sigma_{1/2} \star \mathcal{K}_1 - \mathcal{R}_1 \star \sigma_{1/2} \star \mathcal{K}_2 \\ & - \ln \tau_1^2 \star \sigma_{1/2} \star \mathcal{K}_2 + 2r_1 \star \sigma_{1/2} \star \delta \mathcal{K}_2 + 2r_1 \star s_{1/2} \star \mathcal{K}_1 - 2H \star s_{1/2} \star \mathcal{K}_2. \end{aligned} \quad (2.33)$$

The identity (2.30) was used to simplify the last term.

3. Energy formula and asymptotic form of the identity

One of the important applications of our method is the calculation of the infinite sum

$$\mathcal{E} = -\frac{1}{2\pi} \sum_{Q=1}^{\infty} \int_{-\infty}^{\infty} du L_Q(u) \frac{d\tilde{p}^Q}{du} \quad (3.1)$$

occurring in the expression for the energy of the state [12]. We can apply our formalism to calculate this sum with

$$\mathcal{K}_Q(u, v) \Rightarrow -\frac{1}{2\pi} \frac{d\tilde{p}^Q}{du} = \frac{g}{2\pi} \left[x^{[Q]'}(u) - x^{[-Q]'}(u) \right]. \quad (3.2)$$

This expression satisfies the kernel identity (2.13), but approaches constant limits⁵ for large $|u|$. This last property of this kernel leads to a problem in applying (2.33) since the functions $\mathcal{R}_1, \mathcal{R}_2$ behave as $\ln |u|$ for large $|u|$ and the corresponding convolutions are divergent.

There are two possible solutions of this problem. First we can use the identity

$$x^{[Q]'}(u) - x^{[-Q]'}(u) = i \int_{|v|>2} dv K_Q(u-v) \frac{v+i\epsilon}{\sqrt{4-(v+i\epsilon)^2}}, \quad (3.3)$$

where

$$K_Q(u) = \frac{Qg}{\pi} \frac{1}{Q^2 + g^2 u^2} \quad (3.4)$$

and use our formalism for $\mathcal{K}_Q(u, v) \Rightarrow K_Q(u-v)$ to calculate

$$t_1 = \sum_{Q=1}^{\infty} L_Q \star K_Q. \quad (3.5)$$

Since K_Q is well-behaved at infinity, all convolutions are now convergent and after having computed $t_1(u)$ with our method we can calculate (3.1) by multiplying it with $\frac{ig}{2\pi} \frac{u}{\sqrt{4-u^2}}$ and integrate the result for $|u| > 2$ (slightly above the real line).

An alternative, second solution (which is also applicable to other cases of interest) is to consider the asymptotic limit⁶ of our functions. The asymptotic limit is obtained from the Bethe Ansatz and these asymptotic solutions satisfy TBA integral equations that can be obtained from (4.1,4.2,4.5) by simply deleting all $(1+Y_Q)$ factors occurring on the right hand side of these equations [14]. In our notation the asymptotic TBA equations become

$$\hat{\mathcal{L}}_m^{(0)} = \hat{r}_m^{(0)} - (\hat{r}_{m+1}^{(0)} + \hat{r}_{m-1}^{(0)}) \star s - \hat{f}_2^{(0)} \delta_{m,1} \quad (3.6)$$

(for $m = 1, 2, \dots$) and

$$\hat{L}_Q^{(0)} = \hat{\mathcal{R}}_Q^{(0)} + 2\hat{\mathcal{L}}_{Q-1}^{(0)} \star s - (\hat{\mathcal{R}}_{Q+1}^{(0)} + \hat{\mathcal{R}}_{Q-1}^{(0)}) \star s + (\hat{L}_{Q+1}^{(0)} + \hat{L}_{Q-1}^{(0)}) \star s - \hat{f}_1^{(0)} \delta_{Q,2} \quad (3.7)$$

(for $Q = 2, 3, \dots$) Since the particle rapidities u_j occurring in τ_1 can be treated simply as parameters, we will not distinguish between the asymptotic and “exact” versions of the function f_1 and will use

$$f_1^{(0)} = f_1 = \ln \tau_1^2 \star s. \quad (3.8)$$

We can now apply the same steps to the asymptotic sums

$$\mathcal{A}^{(0)} = \sum_{Q=2}^{\infty} \hat{L}_Q^{(0)} \star \mathcal{K}_Q \quad \mathcal{B}^{(0)} = \sum_{m=1}^{\infty} \hat{\mathcal{L}}_m^{(0)} \star \bar{\mathcal{K}}_{m+1} \quad (3.9)$$

as we did for their “exact” counterparts. We find that $\mathcal{A}^{(0)}$ cannot be determined in this way (it cancels from the equations) but we obtain instead the following identity:

⁵Note that $x(u) = (u - i\sqrt{4-u^2})/2$.

⁶This will be indicated by an upper index ⁽⁰⁾ on the corresponding function.

$$\begin{aligned}
0 = & \sum_{Q=2}^{\infty} (\mathcal{R}_Q^{(0)} - L_Q^{(0)}) \star (\delta\mathcal{K}_Q + 2s_{1/2} \star \delta\mathcal{K}_Q) + (\mathcal{R}_2^{(0)} - L_2^{(0)}) \star \sigma_{1/2} \star \mathcal{K}_1 \\
& - (\mathcal{R}_1^{(0)} - L_1^{(0)}) \star \sigma_{1/2} \star \mathcal{K}_2 - \ln \tau_1^2 \star \sigma_{1/2} \star \mathcal{K}_2 \\
& + 2 \sum_{m=1}^{\infty} r_m^{(0)} \star \sigma_{1/2} \star \delta\mathcal{K}_{m+1} + 2r_1^{(0)} \star s_{1/2} \star \mathcal{K}_1 - 2H^{(0)} \hat{\star} s_{1/2} \star \mathcal{K}_2.
\end{aligned} \tag{3.10}$$

We can recognize that the asymptotic limit of the quantity defined by (2.26) appears here and by rearranging some of the terms it can be expressed as

$$\omega^{(0)} = L_2^{(0)} \star s \star \mathcal{K}_1 - L_1^{(0)} \star \mathcal{K}_1 + \sum_{Q=1}^{\infty} L_Q^{(0)} \star \delta\mathcal{K}_Q. \tag{3.11}$$

From this result it is clear that ω , and using (2.25), also Ω is exponentially small in the asymptotic limit, which of course must be true but is not obvious from the result (2.33).

Let us use the formula (3.10) for the calculation of the energy expression. In this case we have $\delta\mathcal{K}_Q = 0$ for $Q \geq 2$ and we note that if we subtract it from (2.33) the $\ln \tau_1^2$ term cancels from the final result and we obtain

$$\begin{aligned}
t_1 = & L_1 \star K_1 + (\mathcal{R}_2 - \mathcal{R}_2^{(0)} + L_2^{(0)}) \star \sigma_{1/2} \star K_1 - (\mathcal{R}_1 - \mathcal{R}_1^{(0)} + L_1^{(0)}) \star \sigma_{1/2} \star K_2 \\
& + 2(r_1 - r_1^{(0)}) \star s_{1/2} \star K_1 - 2(H - H^{(0)}) \hat{\star} s_{1/2} \star K_2.
\end{aligned} \tag{3.12}$$

Here the difference $\mathcal{R}_2 - \mathcal{R}_2^{(0)}$ decays fast enough for $|u| \rightarrow \infty$ and this fact and the similar behavior of the other terms allow us to apply the identity (3.3) backwards, term by term, in this formula. The final expression for the energy sum (3.1) can be written as follows:

$$\begin{aligned}
\mathcal{E} = & L_1 \star \tilde{J}_1 + (\mathcal{R}_2 - \mathcal{R}_2^{(0)} + L_2^{(0)}) \star \sigma_{1/2} \star \tilde{J}_1 - (\mathcal{R}_1 - \mathcal{R}_1^{(0)} + L_1^{(0)}) \star \sigma_{1/2} \star \tilde{J}_2 \\
& + 2(r_1 - r_1^{(0)}) \star s_{1/2} \star \tilde{J}_1 - 2(H - H^{(0)}) \hat{\star} s_{1/2} \star \tilde{J}_2,
\end{aligned} \tag{3.13}$$

where we introduced the notation $\tilde{J}_Q(u) = -\frac{1}{2\pi} \frac{d\tilde{p}^Q}{du}$.

4. The quasi-local TBA equations

In this section, for the sake of completeness, we present the quasi-local form of the $AdS_5 \times S^5$ mirror TBA equations. The form of the first group of the equations follows from the Y-system relations thus they are local and their form is the the same as that of the simplified

version of the equations.

$$Y_{m|vw}^{(\alpha)} = t_{m|vw}^{(\alpha)} \exp \left\{ \ln \left[\frac{(1 + Y_{m+1|vw}^{(\alpha)})(1 + Y_{m-1|vw}^{(\alpha)})}{(1 + Y_{m+1})} \right] \star s \right\}, \quad m \geq 2, \quad (4.1)$$

$$Y_{1|vw}^{(\alpha)} = t_{1|vw}^{(\alpha)} \exp \left\{ \ln \left[\frac{(1 + Y_{2|vw}^{(\alpha)})}{(1 + Y_2)} \right] \star s + \ln \left[\frac{1 - Y_-^{(\alpha)}}{1 - Y_+^{(\alpha)}} \right] \hat{\star} s \right\}, \quad (4.2)$$

$$Y_{m|w}^{(\alpha)} = t_{m|w}^{(\alpha)} \exp \left\{ \ln \left[(1 + Y_{m+1|w}^{(\alpha)})(1 + Y_{m-1|w}^{(\alpha)}) \right] \star s \right\}, \quad m \geq 2, \quad (4.3)$$

$$Y_{1|w}^{(\alpha)} = t_{1|w}^{(\alpha)} \exp \left\{ \ln \left[1 + Y_{2|w}^{(\alpha)} \right] \star s + \ln \left[\frac{1 - \frac{1}{Y_-^{(\alpha)}}}{1 - \frac{1}{Y_+^{(\alpha)}}} \right] \hat{\star} s \right\}, \quad (4.4)$$

$$Y_Q = t_Q \exp \left\{ \ln \left[\frac{Y_{Q+1} Y_{Q-1} (1 + Y_{Q-1|vw}^{(+)})(1 + Y_{Q-1|vw}^{(-)})}{Y_{Q-1|vw}^{(+)} Y_{Q-1|vw}^{(-)} (1 + Y_{Q+1})(1 + Y_{Q-1})} \right] \star s \right\}, \quad Q \geq 2. \quad (4.5)$$

Here the source terms $t_{\dots}^{(\alpha)}$ correspond to the singularities of the Y -functions within the physical strip (see below). The second group of equations contains the quantity Ω which is a functional of the vector of kernels $\mathcal{K}_Q$. To emphasize this fact from now on Ω in (1.3) will be denoted by $\Omega(\mathcal{K}_Q)$ to make this functional dependence explicit. On the right hand side of the subsequent equations $\Omega(\mathcal{K}_Q)$ always denotes the finite expression obtained for $\Omega(\mathcal{K}_Q)$ in our final formula (2.33). The quasi-local group of equations take the form:

$$\frac{Y_-^{(\alpha)}}{Y_+^{(\alpha)}} = \frac{R_p B_m}{B_p R_m} \exp \{ -L_1 \star K_{1y} - \Omega(K_{Qy}) \}, \quad (4.6)$$

$$\begin{aligned} Y_+^{(\alpha)} Y_-^{(\alpha)} &= \left(t_-^{(\alpha)} \right)^2 \exp \left\{ 2 \ln \left[\frac{1 + Y_{1|vw}^{(\alpha)}}{1 + Y_{1|w}^{(\alpha)}} \right] \star s + L_1 \star [-K_1 + 2K_{xv}^{11} \star s] \right. \\ &\quad \left. - \Omega(K_Q) + 2\Omega(K_{xv}^{Q1} \star s) - \ln \left[\left(\frac{R_p^+ R_p^-}{R_m^+ R_m^-} \right)^2 \frac{Q^{--}}{Q^{++}} \right] \star s \right\}, \end{aligned} \quad (4.7)$$

$$\begin{aligned} \ln Y_1 &= -L\tilde{\mathcal{E}}_1 + f_1 + 2r_1 \star s \hat{\star} K_{y1} \\ &\quad - 2 \ln \left[\frac{1 - Y_-}{1 - Y_+} \right] \hat{\star} s \star K_{vwx}^{11} + 2\mathcal{L}_- \hat{\star} K_-^{y1} + 2\mathcal{L}_+ \hat{\star} K_+^{y1} \\ &\quad + L_1 \star K_{\text{sl}(2)}^{11} + \Omega(K_{\text{sl}(2)}^{Q1}) + 2\Omega(s \star K_{vwx}^{Q-11}), \end{aligned} \quad (4.8)$$

where

$$f_1(u) = - \sum_{j=1}^N \ln S_{\text{sl}(2)}^{1*1}(u_j, u) - 2 [T^\epsilon \star K_{vwx}^{11}] (u)$$

with

$$T^{\pm\epsilon}(u) = \sum_{j=1}^N \ln t(u - u_j \pm i\epsilon), \quad t(u) = \tanh \left(\frac{\pi g u}{4} \right). \quad (4.9)$$

The source terms appearing in the TBA equations (4.1-4.8) contain the objects that are subject to the quantization conditions. In the $\mathfrak{sl}(2)$ sector they can be expressed in terms of the $\tau_m(u) = \prod_{j=1}^{\mathcal{N}_m} t(u - \xi_{m,j})$ and $\tilde{\tau}_m(u) = \prod_{j=1}^{\tilde{\mathcal{N}}_m} t(u - \tilde{\xi}_{m,j})$ functions as follows:

$$t_{m|vw} = \tau_m \tau_{m+2}, \quad m = 1, 2, \dots, \quad (4.10)$$

$$t_{m|w} = \tilde{\tau}_m \tilde{\tau}_{m+2}, \quad m = 1, 2, \dots, \quad (4.11)$$

$$t_Q = \tau_Q^2, \quad Q = 2, 3, \dots, \quad (4.12)$$

$$t_1 = 1, \quad (4.13)$$

$$t_- = \tau_2 / \tilde{\tau}_2. \quad (4.14)$$

Here the $\xi_{m,j}$ and $\tilde{\xi}_{m,j}$ are the objects to be determined by the quantization conditions:

$$1 + Y_{m|vw}^\pm(\xi_{m+1,j}) = 0, \quad m = 1, 2, \dots, \quad j = 1, \dots, \mathcal{N}_{m+1}, \quad (4.15)$$

$$1 + Y_{m|w}^\pm(\tilde{\xi}_{m+1,j}) = 0, \quad m = 1, 2, \dots, \quad j = 1, \dots, \tilde{\mathcal{N}}_{m+1}. \quad (4.16)$$

The set $\{\xi_{1,j}\}_{j=1,\dots,\mathcal{N}_1}$ is equal to the set of physical rapidities $\{u_j\}_{j=1,\dots,N}$ determined by the exact Bethe equations:

$$1 + Y_{1(*)}(u_j) = 0, \quad j = 1, \dots, N \quad (4.17)$$

and the set $\{\tilde{\xi}_{1,j}\}_{j=1,\dots,\tilde{\mathcal{N}}_1}$ is empty by definition. Here $Y_{1(*)}$ denotes Y_1 analytically continued to the physical sheet. For a detailed explanation see refs. [12] and [16]. We note that because of the left-right symmetry of the $\mathfrak{sl}(2)$ sector we have omitted the wing index (α) in the formulae above.

In order to be able to use our result (2.33) in the equations (4.1-4.8) the vector $\delta\mathcal{K}_Q$ associated to the kernels $\mathcal{K}_Q$ appearing in the mirror TBA problem must be known. Since in the expression (2.33) for $\Omega(\mathcal{K}_Q)$, $\delta\mathcal{K}_Q$ with $Q \geq 2$ appears only, we list the results for those $\delta\mathcal{K}_Q$ entering the mirror TBA equations (4.1-4.8) for $Q \geq 2$. They are given by:

$$\delta K_Q = 0, \quad \delta K_{Qy} = 0, \quad \delta K_{xv}^{Q1} = 0, \quad Q \geq 2, \quad (4.18)$$

$$\delta(s \star K_{vwx}^{Q-1,1}) = \delta_{Q,2} s \star s \hat{\star} K_{y1}, \quad \delta K_{\mathfrak{sl}(2)}^{Q1} = -\delta_{Q,2} s, \quad Q \geq 2. \quad (4.19)$$

Using (4.18-4.19) and (2.33) all the $\Omega(\mathcal{K}_Q)$ terms can be explicitly evaluated. The substitution of these expressions into the equations (4.1-4.8) gives the quasi-local form of the mirror TBA equations.

5. Conclusion

In this paper we worked out a method that enabled us to remove all infinite sums from the mirror TBA equations and the energy formula of AdS/CFT and obtained a quasi-local formulation of the TBA problem. Quasi-locality means that in this form at most next to nearest neighbor⁷ Y-functions are coupled by the TBA equations.

⁷with respect to the Y-system diagram of AdS/CFT

We think that the quasi-local formulation of the mirror TBA equations is an important step towards the NLIE formulation of the AdS/CFT spectral problem since using it the application of techniques worked out for the relativistically invariant models [32, 33, 34, 35, 36] becomes possible.

Indeed, the locality of the simplified version of the TBA equations made it possible to find a hybrid-NLIE formulation [37] of the AdS/CFT spectral problem, where the two $SU(2)$ wings of the mirror TBA equations were resummed reducing remarkably the number of unknowns.

Acknowledgements

This investigation was supported by the Hungarian National Science Fund OTKA (under K 77400).

References

- [1] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” Adv. Theor. Math. Phys. **2** (1998) 231 [Int. J. Theor. Phys. **38** (1999) 1113], [hep-th/9711200];
S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” Phys. Lett. B **428** (1998) 105, [hep-th/9802109];
E. Witten, “Anti-de Sitter space and holography,” Adv. Theor. Math. Phys. **2** (1998) 253, [hep-th/9802150].
- [2] G. Arutyunov and S. Frolov, “Foundations of the $AdS_5 \times S^5$ Superstring. Part I,” J. Phys. A **42** (2009) 254003 [arXiv:0901.4937 [hep-th]].
- [3] J. Ambjorn, R. A. Janik and C. Kristjansen, “Wrapping interactions and a new source of corrections to the spin-chain / string duality,” Nucl. Phys. B **736** (2006) 288 [arXiv:hep-th/0510171].
- [4] G. Arutyunov and S. Frolov, “On String S-matrix, Bound States and TBA,” JHEP **0712** (2007) 024, hep-th/0710.1568.
- [5] G. Arutyunov and S. Frolov, “String hypothesis for the AdS mirror,” JHEP **0903** (2009) 152 [arXiv:0901.1417 [hep-th]].
- [6] D. Bombardelli, D. Fioravanti and R. Tateo, “Thermodynamic Bethe Ansatz for planar AdS/CFT: a proposal,” J. Phys. A **42** (2009) 375401 [arXiv:0902.3930].
- [7] N. Gromov, V. Kazakov, A. Kozak and P. Vieira, “Exact Spectrum of Anomalous Dimensions of Planar $N = 4$ Supersymmetric Yang-Mills Theory: TBA and excited states,” Lett.Math.Phys.91:265-287,2010, [arXiv:0902.4458v3 [hep-th]].
- [8] G. Arutyunov and S. Frolov, “Thermodynamic Bethe Ansatz for the AdS Mirror Model,” JHEP **0905** (2009) 068 [arXiv:0903.0141 [hep-th]].
- [9] G. Arutyunov and S. Frolov, “Simplified TBA equations of the $AdS_5 \times S^5$ mirror model,” JHEP 0911:019,2009, arXiv:0907.2647 [hep-th].
- [10] P. Dorey, R. Tateo, “Excited states by analytic continuation of TBA equations,” Nucl. Phys. B **482**, 639 (1996) [arXiv:hep-th/9607167].

- [11] N. Gromov, V. Kazakov and P. Vieira, “Exact AdS/CFT spectrum: Konishi dimension at any coupling,” *Phys.Rev.Lett.*104:211601,2010, [arXiv:0906.4240 [hep-th]].
- [12] G. Arutyunov, S. Frolov and R. Suzuki, “Exploring the mirror TBA,” *JHEP* 1005:031,2010, [arXiv:0911.2224 [hep-th]].
- [13] J. Balog, Á. Hegedűs, ”The Bajnok-Janik formula and wrapping corrections”, *JHEP* 1009:107,2010, [arXiv:1003.4303 [hep-th]].
- [14] N. Gromov, V. Kazakov and P. Vieira, “Integrability for the Full Spectrum of Planar AdS/CFT,” *Phys.Rev.Lett.*103:131601,2009, arXiv:0901.3753 [hep-th].
- [15] A. Cavaglia, D. Fioravanti, R. Tateo, ”Extended Y-system for the AdS_5/CFT_4 correspondence”, *Nucl.Phys. B* **843**, (2011) 302-343, [arXiv:1005.3016 [hep-th]].
- [16] J. Balog, A. Hegedus, ” $AdS_5 \times S^5$ mirror TBA equations from Y-system and discontinuity relations”, [arXiv:1104.4054[hep-th]]
- [17] F. Fiamberti, A. Santambrogio, C. Sieg and D. Zanon, “Wrapping at four loops in N=4 SYM,” *Phys. Lett. B* **666** (2008) 100 [arXiv:0712.3522 [hep-th]].
- [18] V. N. Velizhanin, “The Four-Loop Konishi in N=4 SYM,” arXiv:0808.3832 [hep-th].
- [19] A. V. Kotikov and L. N. Lipatov, “DGLAP and BFKL Evolution Equations in the $\mathcal{N} = 4$ Supersymmetric Gauge Theory,” *Nucl. Phys. B* **661** (2003) 19 [Erratum-ibid. B **685** (2004) 405] [arXiv:hep-ph/0208220].
- [20] A. V. Kotikov, L. N. Lipatov, A. Rej, M. Staudacher and V. N. Velizhanin, “Dressing and Wrapping,” *J. Stat. Mech.* **0710** (2007) P10003 [arXiv:0704.3586 [hep-th]].
- [21] G. Arutyunov, S. Frolov and R. Suzuki, “Five-loop Konishi from the Mirror TBA”, *JHEP* 1004:069,2010, [arXiv:1002.1711 [hep-th]].
- [22] J. Balog, Á. Hegedűs, ”5-loop Konishi from linearized TBA and the XXX magnet”, *JHEP* 1006:080,2010, [arXiv:1002.4142 [hep-th]].
- [23] R. A. Janik and T. Lukowski, “Wrapping interactions at strong coupling – the giant magnon,” *Phys. Rev. D* **76** (2007) 126008 [arXiv:0708.2208 [hep-th]].
- [24] Z. Bajnok and R. A. Janik, “Four-loop perturbative Konishi from strings and finite size effects for multiparticle states,” *Nucl. Phys. B* **807** (2009) 625 [arXiv:0807.0399 [hep-th]].
- [25] Z. Bajnok, R. A. Janik and T. Lukowski, “Four loop twist two, BFKL, wrapping and strings,” *Nucl. Phys. B* **816** (2009) 376 [arXiv:0811.4448 [hep-th]].
- [26] Z. Bajnok, A. Hegedus, R. A. Janik and T. Lukowski, “Five loop Konishi from AdS/CFT,” *Nucl. Phys. B* **827** (2010), 426-456, arXiv:0906.4062 [hep-th].
- [27] T. Lukowski, A. Rej and V. N. Velizhanin, “Five-Loop Anomalous Dimension of Twist-Two Operators,” *Nucl.Phys.B* **831** (2010), 105-132, arXiv:0912.1624 [hep-th].
- [28] N. Gromov, “Y-system and Quasi-Classical Strings,” *JHEP* 1001:112,2010, [arXiv:0910.3608 [hep-th]].
- [29] N. Gromov, V. Kazakov, Z. Tsuboi, ” $PSU(2,2|4)$ character of quasiclasical AdS/CFT” *JHEP* 1007:097,2010, [arXiv:1002.3981 [hep-th]].
- [30] S. Frolov, ”Konishi operator at intermediate coupling”, *J.Phys.A***44**:065401,2011, [arXiv:1006.5032 [hep-th]].

- [31] N. Gromov, D. Serban, I. Shenderovich and D. Volin, "Quantum folded string and integrability: from finite size effects to Konishi dimension", [arXiv:1102.1040[hep-th]], R. Roiban and A. A. Tseytlin, "Semiclassical string computation of strong-coupling corrections to dimensions of operators in Konishi multiplet", [arXiv:1102.1209 [hep-th]] B. C. Vallilo and L. Mazzucato, "The Konishi multiplet at strong coupling", [arXiv:1102.1219 [hep-th]].
- [32] C. Destri, H.J. de Vega, "New thermodynamic Bethe ansatz equations without strings", Phys. Rev. Lett. 69 (1992) 2313-2317;
"Unified approach to thermodynamic Bethe Ansatz and finite size corrections for lattice models and field theories", Nucl. Phys. B438 (1995) 413-454, [arXiv:hep-th/9407117].
- [33] C. Dunning, "Finite size effects and the supersymmetric sine-Gordon models", J.Phys.A36:5463-5476, (2003), [arXiv:hep-th/0210225].
- [34] A. Hegedus, "Finite size effects in the SS-model: two component nonlinear integral equations", Nucl.Phys. B679 (2004) 545-567, [arXiv:hep-th/0310051].
- [35] N. Gromov, V. Kazakov, P. Vieira, "Finite Volume Spectrum of 2D Field Theories from Hirota Dynamics", JHEP 0912:060, (2009), [arXiv:0812.5091 [hep-th]].
- [36] V. Kazakov, S. Leurent, "Finite Size Spectrum of SU(N) Principal Chiral Field from Discrete Hirota Dynamics", [arXiv:1007.1770 [hep-th]]
- [37] R. Suzuki "Hybrid NLIE for the Mirror $AdS_5 \times S^5$ ", [arXiv:1101.5165 [hep-th]].